

AD-A064 114

IRT CORP SAN DIEGO CALIF  
BEAM STUDIES OF MOLECULAR PROCESSES ON METAL SURFACES.(U)  
JAN 79 R L PALMER  
IRT-8162-001

FLD 1

N00014-77-C-0255

NL


UNCLASSIFIED

| OF |

AD  
A064114



END  
DATE  
FILMED  
4-79  
DDC



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

AD A064114

DDC FILE COPY

OFFICE OF NAVAL RESEARCH

Contract N00014-77-C-0255

Task No. 056-527

TECHNICAL REPORT NO. 6

Beam Studies of Molecular  
Processes on Metal Surfaces

by

Robert L. Palmer

Prepared for Publication

in the

Journal of Vacuum Science and Technology

IRT Corporation  
P. O. Box 80817  
San Diego, California 92138

January 22, 1979

Reproduction in whole or in part is permitted for  
any purpose of the United States Government

This document has been approved for public release  
and sale; its distribution is unlimited

12

LEVEL

14 IRT-8162-APP  
TR-6

12 15p

DDC  
RECEIVED  
FEB 2 1979

A

409 388

JOB

79 01 31 025

## UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Beam Studies of Molecular Processes on Metal Surfaces		5. TYPE OF REPORT & PERIOD COVERED Technical Report No. 6✓
7. AUTHOR(s) Robert L. Palmer		6. PERFORMING ORG. REPORT NUMBER IRT 8162-001✓
9. PERFORMING ORGANIZATION NAME AND ADDRESS IRT Corporation✓ P. O. Box 80817 San Diego, California 92138		8. CONTRACT OR GRANT NUMBER(s) N00014-77-C-0255✓
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research, Department of the Navy, 800 N. Quincy Street Arlington, Virginia 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Task No. NR 056-527
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE January 22, 1979
		13. NUMBER OF PAGES
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  This document has been approved for public release and sale; its distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The isosteric heats of adsorption of NO and O <sub>2</sub> have been studied on epitaxially grown Pt(111) using elastic helium scattering as the surface coverage probe. The measured isosteric heats for NO and O <sub>2</sub> are 21 and 26 kcal/mole, respectively. The data also indicate a pre-exponential factor for NO desorption of less than $2 \times 10^7$ . ← 10 to the 7th power		

79 01 31 025

## Beam Studies of Molecular Processes on Metal Surfaces

R. L. Palmer  
IRT Corporation  
P. O. Box 80817  
San Diego, California 92138

ACCESSION NO.	
RTG	DATE
SEC	DATE
CLASSIFICATION	
INITIALS	
BY	
APPROVED FOR PUBLICATION	
DATE	REVIEW OF SPECIAL
A	

Studies of molecular processes on surfaces are motivated by a variety of interests in interface phenomena including catalysis, corrosion and high altitude drag coefficients. Molecular beam studies of gas-surface interactions have been particularly fruitful for probing the details of molecular processes such as adsorption, desorption, energy transfer and chemical reaction. When an atom or molecule collides with a surface it can interact in a number of ways. It may be reflected elastically, partially or completely accommodated or physi- or chemisorbed. Among other things, the energy transfer to the surface varies from zero in the case of elastic reflection to 100 percent for complete accommodation and adsorption.

Since adsorption is the first step in heterogeneous catalysis while accommodation coefficients (energy transfer) are of interest in aerodynamics, a study of the parameters that affect the rate of adsorption of gases on metal surfaces is of particular interest in both of these seemingly unrelated areas. We have used molecular beams to study the adsorption of gases such as  $O_2$ ,  $N_2$ ,  $H_2$ ,  $CH_4$ ,  $C_2H_x$ ,  $NH_3$ ,  $CO$ ,  $CO_2$  and others on both single-crystal and polycrystalline surfaces of metals such as Ag, Au, Ni, Fe, Co, Pd, Pt, and Mg. We have found that for some gas/surface systems the adsorption probability (sticking coefficient  $S$ ), is strongly affected by the physical structure of the surface. Generally speaking, clean, single-crystal surfaces have the lowest sticking probabilities, while rough, polycrystalline surfaces are more likely to partially or fully accommodate or adsorb incident molecules. While this may be due, in part, to multiple collisions on microscopically rough surfaces, subtle changes in the electronic structure of surfaces at step edges and other "low coordination" sites can have a significant influence on the sticking coefficient of molecular species.

Using molecular beams to measure adsorption probabilities as a function of incident angle and kinetic energy, some idea of the gas/surface interaction potential can be obtained. In some cases the adsorption step has an activation barrier that can be overcome most effectively by translational energy while, for others, the barrier is most effectively overcome by internal (vibrational) energy. The importance of this information lies in the ability to modify the surface in such a way as to increase or decrease the activation barrier and thereby vary the sticking coefficient to suit the particular circumstance. The electronic structure is influenced by both the elemental composition and physical structure of

the surface. Sticking coefficients can be lowered by as much as an order of magnitude by eliminating surface structures such as steps, grain boundaries, etc.

The molecular beam technique has long been recognized as the most direct way to study the interaction of molecular and atomic species. Many important studies of the chemical interaction of atoms, molecules, radicals and charged species have been carried out in the gas phase using the beam method. Eastermann and Stern first applied the beam technique to the study of atomic interactions on surfaces some fifty years ago. Their diffraction results were quite significant because they demonstrated the wave properties of particles at a time when de Broglie's hypothesis was a revolutionary concept in physics. However, the value of this and subsequent molecular beam scattering studies on surfaces suffered from the lack of any real ability to characterize the actual surface where the scattering event occurred. Even cleaved crystals such as those used by Eastermann, et al. were subject to adsorption of gases from the vacuum ambient. This was assumed to occur rapidly in the vacuums used in these early experiments since no diagnostic techniques were available for measuring the chemical and physical nature of the vacuum/surface boundary. Before 1964, attempts to study molecular scattering from metal surfaces were largely unsuccessful. Neither diffraction, nor other features one would expect when scattering atoms from smooth, single-crystal metal surfaces were observed in experiments up to that time.


In 1964, Joe N. Smith, Jr. and Howard Saltsburg, working at General Atomic in San Diego, began a series of experiments in which they scattered molecular beams from single-crystal gold and silver surfaces that were grown epitaxially *in situ*. Even in the absence of ultra-high-vacuum, these surfaces could be maintained clean by growing the crystal continuously during the experiment at a fast rate compared with the surface contamination rate. Their results were quite impressive, showing a high degree of specular scattering as expected from an ideal single-crystal surface and repeatability of the results from crystal to crystal. These results spurred both experimental and theoretical interest in gas-surface interactions and considerable fruitful work has subsequently been carried out on a wide variety of metal surfaces. Carefully prepared surfaces of bulk crystals as well as deposited single-crystal films have both been used to give results that appear to approach the ideal, contaminant and defect-free surface required for studies of molecular interactions with surfaces.

This ability to produce well-characterized surfaces led, very naturally, to broadening the scope of molecular beam studies to include the investigation of chemical as well as physical interactions. A number of studies of surface catalyzed chemical reactions, usually on transition metal surfaces, have been performed using molecular beams. Almost all of this work has been reported during the period from 1970 to the present so that this field is really very new. Because of the fundamental nature of molecular beam studies, the systems studied have been those involving rather simple chemical reactions where one hopes to be able to identify each step in the reaction mechanism, and to determine the respective rate parameters. Catalytic reactions studied to date using molecular beams include  $H_2/D_2$  exchange on Pt(111) (Refs. 1,2), Ni(111) (Ref. 3) and Cu(100) (Ref. 4) and Cu(110) (Ref. 4);  $D_2$ , CO and  $C_2H_x$  oxidations on Pt(111) (Refs. 1,5,6) and Pd(111) (Ref. 7) and  $HCOOH$  decomposition on Pt foil


(Ref. 8). Despite the rather straightforward nature of these reactions, both experimental results and interpretations of the data by researchers in this field have often been in disagreement, especially in the earlier work. Researchers are now more familiar with the techniques involved with conducting beam studies of chemical interactions on surfaces and, in particular, with the interpretation of kinetic measurements using phase-lock methods and modulated beams. In many cases the apparent discrepancies between experiments can be explained in terms of artifacts in the measurement technique. Also, the importance of a very careful analysis of the chemical kinetics with respect to alternative reaction models is now more fully appreciated. Real world chemical kinetics do not always follow the rate laws one expects from a very simplified evaluation of a particular reaction scheme. When a careful evaluation of the boundary conditions assumed in the kinetic analysis of a reaction model is made and compared with the actual conditions of the experiment a different conclusion is often reached than that presented in textbook examples. For example, when a diatomic molecule like  $O_2$ ,  $D_2$ , etc., is adsorbed dissociatively on a surface the equilibrium balance between the gas phase and the surface coverage of species  $A_2$  is usually written

$$p_A(1 - \theta_A)^2 = k\theta_A^2$$

for the reaction



so that the surface coverage,  $\theta_A$ , varies as the half power of the partial pressure,  $p_A$ . This relation is valid only in the absence of a competing surface reaction for the removal of A atoms. If, for example, the surface reaction is a simple one of the form



then the surface coverage of A(s) may vary from half order in  $p_A$  to first order in  $p_A$ , depending on the relative rates of the competing reactions. This makes it impossible to distinguish between alternate reaction mechanisms, e.g., Rideal versus Langmuir, without precise knowledge of the surface coverages and the rate constants of the competing reaction paths in a given experiment.

Other factors that can further complicate the kinetics of real surface/adsorbate systems include coverage dependent heats of adsorption, the effects of "precursor" states in the adsorption/desorption process, and phase changes in the adlayer with coverage.

Since our early work on  $H_2/D_2$  exchange on Ni(111), we have stressed the relationship between adsorption and desorption, i.e., that these processes are just time reversals of the same process so that microscopic reversibility applied in general and detailed balance can be applied when the kinetics of adsorption and

desorption are compared at the same temperatures. Shannabarger has pointed out that because of this relationship between adsorption and desorption, precursor states should play an important role in desorption kinetics in an analogous way to that observed in studies of adsorption using beams and other techniques (Ref. 9). It is now being recognized that precursor states do indeed affect adsorption/desorption kinetics in real systems and that it is likely that the interpretation of flash desorption data as well as reaction kinetics are often significantly affected by these states. Molecular beam studies of adsorption are perhaps the most effective way of studying these precursor intermediates, and the importance of these studies to surface chemistry is even clearer now than when they were first begun. Further studies of this type will be of great help in clarifying results obtained by other experimental techniques such as flash desorption and kinetic measurements of surface catalytic reactions.

The foregoing discussion points up the need for more progress in unifying the various areas of surface chemistry. Molecular beam and other recent experiments in surface chemistry have produced a number of surprises that were not anticipated from our rather simplified understanding of the gas-surface boundary. Although the pioneering concepts established early in this century by Langmuir, Knudsen, and others have provided a good framework for understanding the gas-surface interface, further development in this area is clearly needed. In particular, emphasis should be placed on developing the broader implications of results obtained in specialized areas of study such as molecular beam scattering, flash desorption, field ion microscopy, and chemical reaction kinetics on single-crystal surfaces. Efforts are now underway to codify the results of various well-characterized surface studies with significant improvements in the theory of reaction rates, including chemisorption and catalysis being the likely result.

Recently, we have been using molecular beams to study the chemistry of  $\text{NO}_x$  ( $\text{N}_2\text{O}$ ,  $\text{NO}$ , and  $\text{NO}_2$ ) on Pt(111) surfaces. This work is still in progress, but some of our results can be reported now. While the catalysis of  $\text{NO}_x$  on platinum is related to applied areas such as pollution control in automotive exhausts, our interest is to characterize the interaction of  $\text{NO}_x$  with platinum and other group VIII metals in a rather fundamental way.

One technique that can be applied uniquely to single-crystal surfaces is the use of helium beam scattering to measure adsorbate coverages. The technique depends on the fact that adsorbed molecules scatter helium atoms inelastically, whereas the clean surface reflects some fraction of the helium atoms elastically into a well-defined specular ray. This provides a means of probing the surface of coverage and the technique becomes quantitative if the scattering cross section of the adsorbate can be determined. This method has been used to determine the isosteric heats of adsorption of  $\text{NO}_x$  and  $\text{O}_2$  on Pt(111). Figure 1 gives data obtained for helium scattering from Pt(111) as a function of  $\text{O}_2$  pressure and, hence, oxygen coverage on the surface at several surface temperatures. Replotting this data as oxygen pressure versus reciprocal surface temperature at constant coverage (helium attenuation) we obtain the isosteric heat of adsorption. This is done in Figure 3 for 50 percent helium attenuation which is

approximately 20 percent oxygen coverage using appropriate scattering dimensions for oxygen and helium. Similar attenuation data for NO is plotted in Figure 2 and Figure 3 also gives the Clausius-Clapeyron plot for NO, also at 50 percent attenuation.


We first observe that the isosteric heat of adsorption for oxygen is unusually low when compared with other methods that yield values typically in the range 40 to 70 kcal/mole. To understand this result we must consider the method in more detail. Since we rely on inelastic scattering for the detection of adsorbates, the possibility exists that some of the chemisorbed oxygen may not give rise to inelastic scattering, either because it is tightly bound and/or because it lies below the uppermost surface layer and is shielded from scattering with incident helium atoms. A further complication is that oxygen can exist in both atomic and molecular states on platinum and we have no a priori way of determining from helium scattering alone which state we are probing. We assume from the measured slope (26 kcal/mole) that we are dealing with a chemisorbed state and, hence, probably oxygen atoms although this is by no means unequivocal. One more factor to consider is that the effective scattering cross section of loosely bound species may be significantly enhanced because of their greater mobility on the surface.

Considering these factors, we are most likely probing a rather loosely bound oxygen species on Pt(111) (either O<sub>2</sub> or O atoms) at high total coverage and tightly bound oxygen atoms lying below the topmost surface layer are shielded from scattering by helium atoms and not detected.

The chemisorption of NO data of Figure 2 indicates an isosteric heat of adsorption of 21 kcal/mole. Surprisingly, for nondissociative adsorption, the pre-exponential of the desorption rate can be calculated from the data to be

$$S\nu_0 = 2 \times 10^7$$

where S is the initial sticking coefficient for NO on Pt(111). Since  $S \leq 1$ , the pre-exponential is quite low, in the range normally expected for dissociative adsorption where desorption involves a concerted process. We speculate at this point that chemisorbed NO, while perhaps not completely dissociated, is bound to the surface at both the nitrogen and oxygen atoms, i.e., lying down rather than erect as in the case of CO adsorption. The chemisorption of NO<sub>2</sub> is consistent with the process



Strenuous attempts were made to react all three species, N<sub>2</sub>O, NO, and NO<sub>2</sub> with CO without success. No CO<sub>2</sub> reaction product could be observed by the molecular beam technique. The upper limit of reaction probability is less than 10<sup>-3</sup> per CO collision with the target surface. While this does not preclude the detection of reaction by other more sensitive techniques, the result is surprising for the case of CO + NO<sub>2</sub> since CO reacts rapidly with surface oxygen

atoms on Pt(111). Evidently the presence of NO(s) inhibits the reaction between O(s) and CO(s). Experiments are planned to study the reaction  $\text{CO} + \text{O}_2 + \text{NO}$  to verify the inhibition effect of NO on the oxygen rate of CO.

#### REFERENCES

1. J. N. Smith, Jr. and R. L. Palmer, J. Chem. Phys. 56, 13 (1972).
2. S. L. Bernasek and G. A. Somorjai, J. Chem. Phys. 62, 3149 (1975).
3. R. L. Palmer, J. N. Smith, Jr., H. Saltsburg, and D. R. O'Keefe, J. Chem. Phys. 53, 1666 (1970).
4. M. Balloch, M. J. Cardillo, D. R. Miller, and R. E. Stickney, Surf. Sci. 46, 358 (1974).
5. R. L. Palmer and J. N. Smith, Jr., J. Chem. Phys. 60, 1453 (1974).
6. R. L. Palmer, J. Vac. Sci. Technol. 12, 1403 (1975).
7. T. Engel and G. Ertl, J. Chem. Phys. 69, 1267 (1978).
8. S. C. Dahlberg, G. A. Fisk, and R. R. Rye, J. Catal. 36, 224 (1975).
9. M. R. Shannabarger, Surf. Sci. 64, 43 (1974).




Figure 1




Figure 2




Figure 3

# TECHNICAL REPORT DISTRIBUTION LIST

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Arlington, Virginia 22217 Attn: Code 472	2	Defense Documentation Center Building 5, Cameron Station Alexandria, Virginia 22314	12
Office of Naval Research Arlington, Virginia 22217 Attn: Code 102IP 1	6	U.S. Army Research Office P.O. Box 12211 Research Triangle Park, N.C. 27709 Attn: CRD-AA-IP	1
ONR Branch Office 536 S. Clark Street Chicago, Illinois 60605 Attn: Dr. Jerry Smith	1	Naval Ocean Systems Center San Diego, California 92152 Attn: Mr. Joe McCartney	1
ONR Branch Office 715 Broadway New York, New York 10003 Attn: Scientific Dept.	1	Naval Weapons Center China Lake, California 93555 Attn: Head, Chemistry Division	1
ONR Branch Office 1030 East Green Street Pasadena, California 91106 Attn: Dr. R. J. Marcus	1	Naval Civil Engineering Laboratory Port Hueneme, California 93041 Attn: Mr. W. S. Haynes	1
ONR Branch Office 760 Market Street, Rm. 447 San Francisco, California 94102 Attn: Dr. P. A. Miller	1	Professor O. Heinz Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
ONR Branch Office 495 Summer Street Boston, Massachusetts 02210 Attn: Dr. L. H. Peebles	1	Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
Director, Naval Research Laboratory Washington, D.C. 20390 Attn: Code 6100	1	Office of Naval Research Arlington, Virginia 22217 Attn: Dr. Richard S. Miller	1
The Asst. Secretary of the Navy (R&D) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1		
Commander, Naval Air Systems Command Department of the Navy Washington, D.C. 20360 Attn: Code 310C (H. Rosenwasser)	1		

# TECHNICAL REPORT DISTRIBUTION LIST

	<u>No. Copies</u>		<u>No. Copies</u>
<del>Dr. D. A. Vroom</del>		Dr. R. W. Vaughan	
<del>IRT</del>		California Institute of Technology	
<del>P.O. Box 80817</del>		Division of Chemistry & Chemical	
<del>San Diego, California 92138</del>	1	Engineering	
		Pasadena, California 91125	1
Dr. G. A. Somorjai		Dr. Keith H. Johnson	
University of California		Massachusetts Institute of Technology	
Department of Chemistry		Department of Metallurgy and Materials	
Berkeley, California 94720	1	Science	
		Cambridge, Massachusetts 02139	1
Dr. L. N. Jarvis		Dr. M. S. Wrighton	
Surface Chemistry Division		Massachusetts Institute of Technology	
4555 Overlook Avenue, S.W.		Department of Chemistry	
Washington, D.C. 20375	1	Cambridge, Massachusetts 02139	1
Dr. W. M. Risen, Jr.		Dr. J. E. Demuth	
Brown University		IBM Corp.	
Department of Chemistry		Thomas J. Watson Research Center	
Providence, Rhode Island 02912	1	P.O. Box 218	
		Yorktown Heights, New York 10598	1
Dr. M. H. Chisholm		Dr. C. P. Flynn	
Princeton University		University of Illinois	
Chemistry Department		Department of Physics	
Princeton, New Jersey 08540	1	Urbana, Illinois 61801	1
Dr. J. B. Hudson		Dr. W. Kohn	
Rensselaer Polytechnic Institute		University of California (San Diego)	
Materials Division		Department of Physics	
Troy, New York 12181	1	La Jolla, California 92037	1
Dr. John T. Yates		Dr. R. L. Park	
National Bureau of Standards		Director, Center of Materials Research	
Department of Commerce		University of Maryland	
Surface Chemistry Section		College Park, Maryland 20742	1
Washington, D.C. 20234	1		
Dr. Theodore E. Madey			
Department of Commerce			
National Bureau of Standards			
Surface Chemistry Section			
Washington, D.C. 20234	1		
Dr. J. M. White			
University of Texas			
Department of Chemistry			
Austin, Texas 78712	1		

29

TECHNICAL REPORT DISTRIBUTION LIST

<u>No. Copies</u>		<u>No. Cop</u>
	Dr. W. T. Peria Electrical Engineering Department University of Minnesota Minneapolis, Minnesota 55455	1
	Dr. Narkis Tzoar City University of New York Convent Avenue at 138th Street New York, New York 10031	1
	Dr. Chia-wei Woo Northwestern University Department of Physics Evanston, Illinois 60201	1
	Dr. D. C. Mattis Yeshiva University Physics Department Amsterdam Avenue & 185th Street New York, New York 10033	1
	Dr. Robert M. Hexter University of Minnesota Department of Chemistry Minneapolis, Minnesota 55455	1
	Dr. Leonard Wharton James Franck Institute Department of Chemistry 5640 Ellis Avenue Chicago, Illinois 60637	1
	Dr. M. G. Lagally Department of Metallurgical and Mining Engineering University of Wisconsin Madison, Wisconsin 53706	1
	Dr. Robert Gomer James Franck Institute Department of Chemistry 5640 Ellis Avenue Chicago, Illinois 60637	1
	Dr. R. F. Wallis University of California (Irvine) Department of Physics Irvine, California 92664	1